

International Journal of Radiology and Diagnostic Imaging



E-ISSN: 2664-4444
P-ISSN: 2664-4436
www.radiologypaper.com
IJRDI 2022; 5(4): 18-24
Received: 12-08-2022
Accepted: 15-09-2022

Dr. Ravi N

Professor, Department of
radio-diagnosis, Bangalore
Medical College and Research
Institute Bangalore,
Karnataka, India

Dr. Kavya

Postgraduate Student,
Department of Radio-
Diagnosis, Bangalore Medical
College and Research Institute
Bangalore, Karnataka, India

Dr. Darshan R Malkhedkar

Postgraduate Student,
Department of Radio-
Diagnosis, Bangalore Medical
College and Research Institute
Bangalore, Karnataka, India

Dr. Mohit Mouna R

Postgraduate Student,
Department of Radio-
Diagnosis, Bangalore Medical
College and Research Institute
Bangalore, Karnataka, India

Corresponding Author:

Dr. Kavya

Postgraduate student,
Department of radio-diagnosis,
Bangalore medical college and
research institute Bangalore,
Karnataka, India

Spectrum of computed tomography (Ct)-brain findings in COVID-19 patients with CNS manifestations

Dr. Ravi N, Dr. Kavya, Dr. Darshan R Malkhedkar and Dr. Mohit Mouna R

DOI: <http://dx.doi.org/10.33545/26644436.2022.v5.i4a.281>

Abstract

Background: Infection with Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-COV-2) results in Corona Virus Disease 2019 (COVID -19) predominantly it affects respiratory system; however it has become apparent that many other organ systems including central nervous system can also be involved. Imaging plays an essential role in diagnosis of all manifestations of the disease and its complications.

Objective: To study the spectrum of neuro-imaging findings on CT in COVID-19 patients with CNS symptoms.

Material and Methods: Non-contrast CT scan of brain was done in 80 patients of either gender having COVID-19, confirmed by reverse transcriptase polymerase chain reaction (RT-PCR) with CNS manifestations. Images were assessed for presence of abnormal findings like Infarcts, hemorrhage, venous/arterial thrombosis, leukoencephalopathy, white matter disease, hypoxic ischemic injury, encephalitis and posterior reversible encephalopathy syndrome.

Results: A total of 80 patients with COVID-19 were reviewed, out of which CT was abnormal in 35 patients (43.7%). findings included various types of infarcts, intra and extra axial haemorrhages, Diffuse axonal injury and PRES (Posterior Reversible Encephalopathy Syndrome).

Our study demonstrated that the neurological imaging of patients with COVID-19 were variable without a specific pattern but is dominated by ischemic infarcts and intracranial haemorrhages.

Conclusion: COVID-19 is a serious public health crisis and can have neurological manifestations, therefore these patients should be evaluated early for neurological symptoms as timely analysis and management of neurological complications are key to improving the prognosis.

Keywords: Assess refers to process of the critical analysis and valuation and judgement of the status or

Introduction

- Infection with Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-COV-2) results in Corona Virus Disease 2019 (COVID -19), which was declared as a pandemic by the World Health Organization (WHO) on March 11, 2020. COVID-19 affects predominantly the respiratory system, however it has become apparent that many other organ systems including central nervous system (CNS) can also be involved resulting in manifestations like acute cerebro-vascular disease, viral encephalitis and infectious toxic encephalopathy^[1, 2].
- The involvement of multiple systems is due to expression of angiotensin-converting enzyme-2 (ACE-2) receptors in extrapulmonary tissues, which results in entry of SARS-COV-2 and thus viral tissue damage. SARS-COV-2 induced coagulopathy significantly contributes to development of CNS manifestations. Inflammation and endothelial cell dysfunction by the virus are the other potential mechanisms contributing to global brain dysfunction^[1].
- Imaging plays an essential role in the diagnosis of all manifestations of the disease and related complications.

Aims and objectives

- To study the spectrum of neuro-imaging findings on CT in COVID-19 patients with CNS symptoms.

Methods

- This cross-sectional study is done from 01/02/2021 to 01/06/2021, Patients of either gender having COVID-19 confirmed by reverse transcriptase - polymerase chain reaction (RT-PCR) with CNS manifestations like confusion/delirium, headache, seizure, dizziness, neuralgia, ataxia, focal neurological deficit, anxiety or depression were included in this study. Computed tomography (CT) of brain was done in these patients in supine position (A total of 80 patients were included in this study).

Inclusion criteria

- COVID-19 confirmed patients of > 18years age with CNS manifestations.
- Patient with proper consent.

Exclusion criteria

- Pregnant patients.
- Patients with any previous history of brain stroke/ intracranial pathology / neurological disorder/head trauma
- Patients who refused to give consent for the study.

Methodology

Clinical details of the patient, relevant history and clinical examinations findings were recorded and consent for the study was taken.

Technique

CT brain was performed in the patients in supine position and CT images were assessed for Spectrum of neuro-imaging findings like:

Infarct

- Acute infarcts
- Subacute infarcts
- Watershed zone
- Lacunar infarcts

Haemorrhage

Extra-axial haemorrhage

- Intraventricular haemorrhage.
- Subarachnoid haemorrhage.
- Subdural haemorrhage.
- Epidural haemorrhage.

Intra-axial haemorrhage

- Parenchymal /lobar haemorrhage
- Striato-thalamic haemorrhage
- Brainstem haemorrhage
- Cerebellar haemorrhage

Micro-haemorrhage

- Cerebral venous thrombosis / Thrombus in intracranial arteries
- Leukoencephalopathy/White matter disease / Diffuse axonal injury
- Hypoxic ischemic encephalopathy
- PRES (Posterior Reversible Encephalopathy Syndrome)
- Encephalitis.

Statistical Analysis

The data collected will be entered in Microsoft excel and analyzed by SPSS version 24.0

Data analyzed by descriptive statistics namely percentage of proportion, graphs, tables and images whatever applicable.

Results

A total of 80 patients with COVID-19 were reviewed, out of which CT was abnormal in 35 patients (43.7%). findings included various types of infarcts, intra and extra axial haemorrhages, Diffuse axonal injury and PRES (Posterior Reversible Encephalopathy Syndrome).

Our study demonstrated that the neurological imaging of patients with COVID-19 were variable without a specific pattern but is dominated by ischemic infarcts and intracranial haemorrhages.

Patient's demographic parameters

Table 1: Sex distribution of SBO

Characteristics	Number (N)	Percentage(P)
Sex		
Male	49	61.2%
Female	31	38.8%

Table 2: Age distribution of SBO

Characteristics	Number (N)	Percentage(P)
Age		
< 50 years	19	23.8%
> 50 years	61	76.2%

Table 3: Indications for the CT brain in COVID-19 patients

Indications	Number (N)	Percentage(%)
Delirium/confusion	30	37.5 %
Focal neurological deficit	13	16.2 %
Anxiety / depression	11	13.8%
Headache	9	11.2%
Dizziness	7	8.8%
Seizure	5	6.2%
Neuralgia	3	3.8%
Ataxia	2	2.5%




Fig 1: PIE Chart depicting gender distribution of SBO





Fig 2: PIE Chart depicting Age distribution of SBO

Table 4: Distribution of various CT brain findings in COVID-19 patients

Ct brain findings:	Number (N)	Percentage (%)
Infarcts	21	60%
- Acute Infarcts	7	(20%)
- Subacute infarcts	6	(17.1%)
- watershed	3	(8.5%)
- Lacunar	5	(14.2%)
Hemorrhages	9	28.5%
Intra-axial Hemorrhages	7	(20%)
- Lobar	4	(11.4%)
- Striato-thalamic	1	(2.8%)
- Micro-hemorrhages	1	(2.8%)
-Brain-stem and cerebellum	1	(2.8%)
Extra-axial Hemorrhages	2	(5.7%)
- Intraventricular	1	(2.8%)
- Subarachnoid	1	(2.8%)
Cerebral venous thrombosis	3	8.5%
-superior sagittal sinus and cortical veins	1	(2.8%)
-Transverse sinus	2	(5.7%)
PRES (Posterior Reversible Encephalopathy Syndrome)	1	2.8%
Diffuse axonal injury	1	2.8%

Representative Images Infarcts



Acute infarct

Sub-acute infarct

Watershed infarct

Lacunar infarct

Fig 3: CT images of various types of infarcts in covid-19 patients.

Hemorrhages

Intra-axial Hemorrhages




Fig 4: Extra-axial Hemorrhages




Fig 5: CT images of various types of hemorrhages in covid-19 patients.

Cerebral venous thrombosis




Fig 6: CT images of cerebral thrombosis in covid-19 patients.



Fig 7: CT image of diffuse axonal injury in covid-19 patient.




Fig 8: CT images of PRES (vasogenic edema in parieto-occipital region) in covid-19 patient.

Discussion

Infection with Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-COV-2) results in Corona Virus Disease 2019 (COVID -19), it affects predominantly the respiratory system, however it has become apparent that many other organ systems including central nervous system (CNS). The neurological manifestations are hypothesized to be caused by multiple factors such as damage to specific receptors, secondary hypoxia, cytokine-related injury and retrograde travel along the olfactory nerve and bulb.

Additionally global inflammatory markers such as interleukin (IL)-6, (IL)-12, (IL)-15 and tumor necrosis factor alpha can activate glial cells and produce inflammatory responses. Cytokine storm syndrome and secondary hemophagocytic lympho-histiocytosis can occur in a sub-group of patients leading to hyperinflammatory syndrome.

These systemic consequences along with alveolar damage cause severe hypoxia which leads to cerebro-vascular vasodilatation, cerebral edema and ischemia.

The occurrence of a diverse mechanism in COVID-19 patients, possibly related to endothelial damage³, resulting in impaired microcirculatory function and vascular fragility with micro-hemorrhages or thrombotic microangiopathy⁴

A unique spike glycoprotein receptor binding domain of SARS -C0V-2 confers affinity of virus for the ACE-2 receptors, virus utilizes the ACE-2 receptor for the entry into the host cells. These receptors are also expressed in endothelial cells of blood-brain barrier, therefore patients with COVID-19 may be at risk of thrombogenesis and cerebral ischemia due to both biochemical hypercoagulable states and direct vascular endothelial injury.




Fig 9: The mechanisms of corona-viruses infections and neurological damage caused by corona virus²




Fig 10: The neurophysiopathogenic mechanisms of SARS-CoV-2².

Conclusion

Corona virus-19 (C0VID-19) is a serious public health crisis and can have neurological manifestations. Therefore, patients with covid-19 infections should be evaluated early for neurological symptoms - timely analysis and management of neurological complications are key to improving the prognosis.

Conflict of Interest

Not available

Financial Support

Not available

References

1. Revzin MV, Raza S, Srivastava NC, Warshawsky R, D'Agostino C, Malhotra A, *et al.* Multisystem Imaging Manifestations of COVID-19, Part 2: From Cardiac Complications to Pediatric Manifestations. *Radiographics*. 2020;40(7):1866-92.
2. Wu Y, Xu X, Chen Z, Duan J, Hashimoto K, Yang L, *et al.* Nervous system involvement after infection with COVID-19 and other coronaviruses. *Brain Behav Immun*. 2020;87:18-22.
3. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, *et al.* Endothelial cell infection and endothelitis in COVID-19. *Lancet*. 2020;395(10234):1417-1418. Doi:10.1016/S0140-6736(20)30937-5.
4. Ciceri F, Beretta L, Scandroglio AM, Colombo S, Landoni G, Ruggeri A, *et al.* Microvascular COVID-19 lung vessels obstructive thromboinflammatory syndrome (MicroCLOTS): an atypical acute respiratory distress syndrome working hypothesis. *Crit care Resusc*. 2020;2:10-38.

How to Cite This Article

Ravi N, Kavya, Malkhedkar DR, Mouna MR. Spectrum of computed tomography (Ct)-brain findings in COVID-19 patients with CNS manifestations. *International Journal of Radiology and Diagnostic Imaging*. 2022;5(4):18-24.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.