

# International Journal of Radiology and Diagnostic Imaging



E-ISSN: 2664-4444  
P-ISSN: 2664-4436  
[www.radiologypaper.com](http://www.radiologypaper.com)  
IJRDI 2020; 3(1): xx-xx  
Received: 20-11-2019  
Accepted: 24-12-2019

**Dr. P Suresh**  
Associate Professor,  
Department of Radiodiagnosis,  
The Oxford Medical College,  
Hospital and Research Centre,  
Yadavanahalli, Attibele Hobli,  
Anekal, Bangalore,  
Karnataka, India

## A study of hemodynamics and complications when radiographic contrast media are used

**Dr. P Suresh**

**DOI:** <http://dx.doi.org/10.33545/26644436.2020.v3.i1c.76>

### Abstract

Visualising soft tissues is always challenging in the field of radiology. Majority of times the task is made easy by using radio-opaque contrast media. They are used to improve the contrast of the internal organs that are usually not visualised. Many articles are published regarding the side effects. The side effects are reported to range from a simple itch to life threatening conditions like contrast medium induced nephropathy. Although rare, its effects cannot be neglected. This study is one such attempt to understand the effects of radiographic contrast media on the hemodynamic factors and also to report the most common complications.

**Keywords:** Complications, Hemodynamics, contrast, radiology

### Introduction

Visualising soft tissues is always challenging in the field of radiology. Majority of times the task is made easy by using radio-opaque contrast media. They are used to improve the contrast of the internal organs that are usually not visualised otherwise. Many articles are published regarding the side effects. The side effects are reported to range from a simple itch to life threatening conditions like contrast medium induced nephropathy<sup>1</sup>. Although rare, its effects cannot be neglected.

The currently used contrast media are based on 2,4,6-tri-iodinated benzene ring and are used mainly for diagnostic as well as therapeutic purposes. Iodine-based contrast media are usually classified as ionic or non-ionic and as monomeric and dimeric and are commonly used to visualize vessels, tissues, organs, and the urinary tract. They are helpful in differentiating between normal and pathological areas. They are usually safe but sometimes it can be very harmful and many have reported that it would be more adverse when the patient has already been sensitised for the same drug before.<sup>2</sup> Hypersensitivity reactions, thyroid dysfunction, and contrast-induced nephropathy are the major adverse effects that can be seen in patients undergoing the contrast study. Skin allergy can be acute or delayed and has been reported for upto 48 hours after the procedure.<sup>3-8</sup>

The contrast medium is rich in iodine and sometimes can result in iodine induced hypothyroidism<sup>9,10</sup>. The mechanisms that leads to Contrast Induced Nephrotoxicity (CIN) have not been fully explained and may be due to several factors. The generally held view is that CIN is caused by a combination of a reduction in medullary blood flow leading to hypoxia and direct renal tubular damage due to toxicity of contrast media upon the kidneys. Hypoxia may lead to the formation of reactive oxygen species (ROS) and it has been argued that these in turn are responsible for contrast media toxicity.<sup>9,10,11</sup>. This study is one such attempt to understand the effects of the contrast media on the hemodynamic factors and also to report the most common complications.

### Aims and Objectives

To study the hemodynamics and complications when radiographic contrast media are used in Computed Tomography (CT) and Radiography (X-Rays).

### Materials and Methods

This study was done in the Department of Radiodiagnosis at The Oxford Medical College, Hospital and Research Centre, Bangalore.

### Corresponding Author:

**Dr. P Suresh**  
Associate Professor,  
Department of Radiodiagnosis,  
The Oxford Medical College,  
Hospital and Research Centre,  
Yadavanahalli, Attibele Hobli,  
Anekal, Bangalore,  
Karnataka, India

The study was done from January 2018 to November 2019. Four Hundred patients were chosen who underwent the contrast Radiography and the CT.

Radiography (X-Rays) and Computed Tomography scans (CT scans) were done after taking all necessary precautions. The patients were divided into four groups. The first group which consisted of 100 patients were normal patients. The second which consisted of 100 patients were known hypertensives. The third group which consisted of 100 patients were known diabetics and the fourth group of 100 patients were aged more than 65 years.

The haemodynamics in terms of the heart rate and blood pressure was monitored at five minutes interval for nine consecutive times. The report was noted and the Anova statistics for heart rate, systolic blood pressure and diastolic blood pressure was done.

The complications were noted and descriptive statistics was done.

**Inclusion Criteria**

Only patients undergoing contrast Radiography and CT scan were chosen for the study.

**Exclusion criteria**

- Patients with known nephropathy or on nephrotoxic drugs.
- Patients with high serum creatinine.

**Results**

**Table 1: Heart Rate**

ANOVA for heart rate		
	F	Significance(p)
intra group difference	103.029	<0.001
Inter group difference	2.775	0.001

There was a significant intragroup ( $p<0.001$ ) and inter group differences in heart rate during the period of study ( $p=0.001$ ). Heart rate variations were statistically significant on comparing group D with group A ( $p<0.001$ ), group B ( $p=0.32$ ) and group C ( $p=0.11$ ) during the observation period.

**Table 2: Intergroup difference in heart rate**

Intergroup difference in heart rate		Significance(p)
Group A	Group B	.147
	Group C	.300
	Group D	<0.001
Group B	Group C	.983
	Group D	.032
Group C	Group D	.011

**Table 3: Systolic Blood Pressure**

ANOVA for systolic BP		
	F	Significance(p)
Intra Group difference	236.83	<0.001
Inter Group difference	6.096	<0.001

There was significant difference in the BP in all groups during the period of study ( $p<0.001$ ) with a significant inter group differences ( $p<0.001$ ). The difference was seen between of the groups A and D ( $p<0.001$ ), B and C ( $p<0.11$ ), B and D ( $p<0.001$ ), and C and D ( $p=0.041$ ).

**Table 4: Inter group difference in systolic BP**

Inter group difference in systolic BP		Significance(p)
Group A	Group B	.915
	Group C	.076
	Group D	<.0001
Group B	Group C	.011
	Group D	<0.001
Group C	Group D	.041

**Table 5: Diastolic Blood Pressure**

ANOVA for diastolic BP		
		Significance (p).
Infra group difference	192.247	<0.001
Inter group difference	5.726	<0.001

There is significant intragroup difference ( $p<0.001$ ) and intergroup difference ( $p<0.001$ ) in diastolic BP during period of study. Significant differences exist between group A and group C ( $p=0.004$ ), group A and group D ( $P<0.001$ ), group B and group D ( $p=0.048$ ), group B and group 0 ( $p<0.001$ ).

**Table 6: Inter group difference in diastolic BP**

GROUP		Significance (p)
Group A	Group B	.843
	Group C	.004
	Group D	<.001
Group B	Group C	.048
	Group D	<.001
Group C	Group D	.247

**Table 7: Complications**

Complications	Group A	Group B	Group C	Group D
Allergic Symptoms	2	1	Nil	1
Abnormal Thyroid function	Nil	Nil	3	Nil
Nephropathy	Nil	1	1	1

**Discussion**

Use of lowest dosage of contrast media must be employed. The development of newer imaging technologies has facilitated faster image acquisition; this has enabled radiologists to perform studies with less intravascular contrast, because the duration of time over which contrast needs to be administered has shortened. Considering that high doses of contrast media are required for percutaneous coronary intervention, several formulas have been suggested to calculate the dosage that is least dangerous for renal function. Cigarroa’s formula suggests the following contrast material limit: 5 mL of contrast per kilogram body weight/serum creatinine (mg/dL) with maximum dose acceptable of 300 mL for diagnostic coronary arteriography. Laskey’s formula suggests the volume of contrast to calculated creatinine clearance ratio with a cut-off point for the ratio at 3.7 for percutaneous coronary intervention: a ratio >3.7 would be associated, following contrast use, with a decrease in creatinine clearance and a significant increase in mortality of patients with ST elevation myocardial infarction. More recently the cut-off point for Laskey’s formula has been placed at 2.0: below a ratio of 2.0 CIN would be a rare complication of percutaneous coronary intervention, but it would increase dramatically at a ratio of 3.0. Some authors have suggested using the ratio of grams of iodine to the calculated creatinine clearance; a ratio 1.42,

or better 1.0, would prevent CIN. But the different results obtained by different authors suggest that this needs to be validated further before being accepted in clinical practice, considering also that patients are not a homogeneous group, since some of them may have complications such as hypotension, shock, and reduced left ventricular systolic function that are themselves risks for CIN.

### Conclusion

Although rare, life threatening complications have been reported with the administration of radio-opaque contrast media for radiological investigations and radiological interventions. Hence adequate precautionary measures have to be taken to ensure the safety of patients undergoing these procedures.

### References

1. Lightfoot CB, Abraham RJ, Mammen T, Abdoell M, Kapur S, Abraham RJ. "Survey of radiologists' knowledge regarding the management of severe contrast material-induced allergic reactions, *Radiology*. 2009; 251(3):691-696,
2. Singh J, Daftary A. Iodinated contrast media and their adverse reactions, *Journal of Nuclear Medicine Technology*. 2008; 36(2):69-76.
3. Thomson KR, Varma DK. Safe use of radiographic contrast media, *Australian Prescriber*. 2010; 33(1):19-22,
4. Loh S, Bagheri S, Katzberg RW, Fung MA, Li C. "Delayed adverse reaction to contrast-enhanced CT: a prospective single-center study comparison to control group without enhancement, *Radiology*, 2010; 255(3):764-771
5. Schild HH, Kuhl CK, Hübner-Steiner U, Böhm I, Speck U. Adverse events after unenhanced and monomeric and dimeric contrast-enhanced CT: a prospective randomized controlled trial, *Radiology*, 2006; 240(1):56-64.
6. Sutton AGC, Finn P, Grech ED *et al.*, Early and late reactions after the use of iopamidol 340, ioxaglate 320, and iodixanol 320 in cardiac catheterization, *American Heart Journal*, 2001; 141(4):677-683,
7. Sutton AGC, Finn P, Campbell PG *et al.*, Early and late reactions following the use of iopamidol 340, iomeprol 350 and iodixanol 320 in cardiac catheterization, *Journal of Invasive Cardiology*. 2003; 15(3):133-138.
8. Sendeski MM. Pathophysiology of renal tissue damage by iodinated contrast media, *Clinical and Experimental Pharmacology and Physiology*. 2011; 38(5):292-299,
9. Van der Molen AJ, Thomsen HS, Morcos SK *et al.*, Effect of iodinated contrast media on thyroid function in adults, *European Radiology*. 2004; 14(5):902-907,
10. Katzberg RW, Haller C. Contrast-induced nephrotoxicity: clinical landscape, *Kidney International*. Supplement. 2006; 69:S3-S7,
11. Moisey RS, McPherson S, Wright M, Orme SM. Thyroiditis and iodide mumps following an angioplasty, *Nephrology Dialysis Transplantation*. 2007; 22(4):1250-1252.
12. Giaccia AJ, Simon MC, Johnson R. The biology of hypoxia: the role of oxygen sensing in development, normal function, and disease, *Genes and Development*, 2004; 18(18):2183-2194.
13. Heyman SN, Rosen S, Khamaisi M, Idée J,

- Rosenberger C. Reactive oxygen species and the pathogenesis of radiocontrast-induced nephropathy, *Investigative Radiology*. 2010; 45(4):188-195.
14. Dawson P, Harrison MJG, Weisblatt E. Effect of contrast media on red cell filtrability and morphology, *British Journal of Radiology*. 1983; 56(670):707-710.